

10/554,176

Connecting via Winsock to STN

Welcome to STN International! Enter x:x



* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 15:36:41 ON 31 JUL 2008

=> file reg

=>

Uploading C:\Program Files\Stnexp\Queries\10554176.str



chain nodes :

17 18

ring nodes :

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16

ring/chain nodes :

19

chain bonds :

8-19 10-14 16-17 17-18

ring bonds :

1-2 1-6 2-3 3-4 4-5 4-7 5-6 5-10 7-8 8-9 9-10 11-12 11-16 12-13 13-14
14-15 15-16

exact/norm bonds :

8-19 16-17 17-18

exact bonds :

10-14

normalized bonds :

1-2 1-6 2-3 3-4 4-5 4-7 5-6 5-10 7-8 8-9 9-10 11-12 11-16 12-13 13-14
14-15 15-16

Match level :

10/554,176


1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom
11:Atom 12:Atom 13:Atom 14:Atom 15:Atom 16:Atom 17:Atom 18:Atom 19:CLASS

L1 STRUCTURE UPLOADED

=> d l1

L1 HAS NO ANSWERS

L1 STR



Structure attributes must be viewed using STN Express query preparation.

=> s l1 full

L3 201 SEA SSS FUL L1

=> file ca

=> s l3

L4 4 L3

=> d ibib abs fhistr 1-4

L4 ANSWER 1 OF 4 CA COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 148:449430 CA


TITLE: Design, synthesis, and biological evaluation of
8-biarylquinolines: A novel class of PDE4 inhibitors
AUTHOR(S): Gallant, Michel; Chauret, Nathalie; Claveau, David;
Day, Stephen; Deschenes, Denis; Dube, Daniel; Huang,
Zheng; Lacombe, Patrick; Laliberte, France; Levesque,

Jean-Francois; Liu, Susana; Macdonald, Dwight;
Mancini, Joseph; Masson, Paul; Mastracchio, Anthony;
Nicholson, Donald; Nicoll-Griffith, Deborah A.;
Perrier, Helene; Salem, Myriam; Styhler, Angela;
Young, Robert N.; Girard, Yves

CORPORATE SOURCE: Department of Medicinal Chemistry, Merck Frosst Center
for Therapeutic Research, Pointe Claire-Dorval, QC,
H9R 4P8, Can.


SOURCE: Bioorganic & Medicinal Chemistry Letters (2008),
18(4), 1407-1412
CODEN: BMCLE8; ISSN: 0960-894X

PUBLISHER: Elsevier Ltd.
DOCUMENT TYPE: Journal
LANGUAGE: English
GI



- AB The structure-activity relationship of a novel series of 8-biarylquinolines, e.g., I, acting as type 4 phosphodiesterase (PDE4) inhibitors is described herein. Prototypical compds. from this series are potent and non-selective inhibitors of the four distinct PDE4 (IC₅₀ < 10 nM) isoenzymes (A-D). In a human whole blood in vitro assay, they inhibit (IC₅₀ < 0.5 μM) the LPS-induced release of the cytokine TNF-α. Optimized inhibitors were evaluated in vivo for efficacy in an ovalbumin-induced bronchoconstriction model in conscious guinea pigs. Their propensity to produce an emetic response was evaluated by performing pharmacokinetic studies in squirrel monkeys. This work has led to the identification of several compds. with excellent in vitro and in vivo profiles, including a good therapeutic window of efficacy over emesis.
- IT 1019332-34-3P
RL: PAC (Pharmacological activity); PKT (Pharmacokinetics); PUR (Purification or recovery); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
(preparation, phosphodiesterase 4 inhibitory activity, and SAR of biarylquinolines)
- RN 1019332-34-3 CA
- CN Cyclopropanecarboxylic acid, 2-[3'-[6-[1-methyl-1-(methylsulfonyl)ethyl]-8-quinolinyl][1,1'-biphenyl]-4-yl]-, (1R,2R)- (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 2 OF 4 CA COPYRIGHT 2008 ACS on STN
 141:410960 CA
 TITLE: Preparation of 8-(3-biaryl)phenylquinoline
 phosphodiesterase-4 inhibitors
 INVENTOR(S): Dube, Daniel; Dube, Laurence; Gallant, Michel;
 Lacombe, Patrick; Deschenes, Denis; MacDonald, Dwight
 PATENT ASSIGNEE(S): Merck Frosst Canada & Co., Can.
 SOURCE: PCI Int. Appl., 129 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------|----------|-----------------|----------|
| WO 2004096220 | A1 | 20041111 | WO 2004-CA622 | 20040427 |
| W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW | | | | |
| RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG | | | | |
| AU 2004234190 | A1 | 20041111 | AU 2004-234190 | 20040427 |
| CA 2523336 | A1 | 20041111 | CA 2004-2523336 | 20040427 |
| EP 1635829 | A1 | 20060322 | EP 2004-729586 | 20040427 |

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
 IE, SI, LT, LV, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK
 CN 1812787 A 20060802 CN 2004-80018346 20040427
 JP 2006524638 T 20061102 JP 2006-504121 20040427
 US 20060223850 A1 20061005 US 2005-554176 20051021
 IN 2005DN04934 A 20070928 IN 2005-DN4934 20051027
 PRIORITY APPLN. INFO.: US 2003-466542P P 20030430
 WO 2004-CA622 W 20040427
 OTHER SOURCE(S): MARPAT 141:410960
 GI


* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The title 8-phenylquinolines I [S1-S3 = H, OH, halo, alkyl, etc.; R1 = CO2aryl, CONHaryl, CONHheteroaryl, etc.; Ar1, Ar2 = (hetero)aryl or an N-oxide thereof; R2 = H, aryl, haloaryl, heterocyclyl, etc.; R3 = H, alkyl, hydroxyalkyl, etc.; R4 = H, halo, CN, alkyl, etc.] which are PDE4 inhibitors, were prepared E.g., a multi-step synthesis of II (no characterization data given for intermediates), which showed IC50 of 0.155 μ M in LPS and FMLP-induced TNF- α and LTB4 assays in human whole blood, was given. The pharmaceutical compns. comprising the compound I are claimed.

IT 791630-50-7P
 RL: PAC (Pharmacological activity); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)
 (preparation of 8-(3-biaryl)phenylquinoline phosphodiesterase-4 inhibitors)

RN 791630-50-7 CA

CN 2-Thiazolamine, 5-[3-[6-[1-methyl-1-(methylsulfonyl)ethyl]-8-quinolinyl]phenyl]-4-[4-(methylsulfonyl)phenyl]- (CA INDEX NAME)



REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 3 OF 4 CA COPYRIGHT 2008 ACS on STN
 ACCESSION NUMBER: 140:59526 CA
 TITLE: Preparation of 8-(biaryl)quinolines as PDE4 inhibitors
 INVENTOR(S): Deschenes, Denis; Dube, Daniel; Dube, Laurence;
 Gallant, Michel; Girard, Yves; Lacombe, Patrick;
 MacDonald, Dwight
 PATENT ASSIGNEE(S): Merck Frosst Canada & Co., Can.
 SOURCE: PCT Int. Appl., 122 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------------------|----------|-----------------|------------|
| WO 2004000814 | A1 | 20031231 | WO 2003-CA957 | 20030623 |
| W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW | | | | |
| RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG | | | | |
| CA 2490043 | A1 | 20031231 | CA 2003-2490043 | 20030623 |
| AU 2003243870 | A1 | 20040106 | AU 2003-243870 | 20030623 |
| EP 1517895 | A1 | 20050330 | EP 2003-760540 | 20030623 |
| EP 1517895 | B1 | 20070314 | | |
| R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK | | | | |
| JP 2006502104 | T | 20060119 | JP 2004-514482 | 20030623 |
| AT 356808 | T | 20070415 | AT 2003-760540 | 20030623 |
| ES 2282667 | T3 | 20071016 | ES 2003-760540 | 20030623 |
| US 20050234238 | A1 | 20051020 | US 2004-517416 | 20041208 |
| US 7153968 | B2 | 20061226 | | |
| PRIORITY APPLN. INFO.: | | | US 2002-391364P | P 20020625 |
| | | | US 2002-428313P | P 20021122 |
| | | | WO 2003-CA957 | W 20030623 |
| OTHER SOURCE(S): | MARPAT 140:59526 | | | |
| GI | | | | |


* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Title compds. I [wherein A = C or N; X = Ph, pyridyl, pyrazinyl, thiaphenyl, quinolinyl, benzofuranlyl, oxadiazolyl, diazolyldipyrindinyl, imidazolyldipyrindinyl, oxadiazolylphenyl, benzodioxolyl; R1 = H, halo, or (un)substituted alkanoyl, cyclo/alkyl, alkenyl; R2, R3 = independently H, halo, OH, CN, NO2, or dialkenyl/dicycloalkyl/alkyl, alkenyl, wide variety

of C-containing and heteroat. groups and/or functional groups optionally linked by C1-4alkyl; R2 optionally forms a double bond with an adjoining bond; R4 = H, halo; any ring nitrogen optionally forms N-oxide and N-chloride; and pharmaceutically acceptable salts thereof] were prepared as phosphodiesterase IV (PDE4) inhibitors. For example, II was prepared by Suzuki cross-coupling of quinoline III with 2-bromo-3-chlorothiophene. One hundred fifty-five invention compds. suppressed PDE4 with IC50 values ranging from 36 μ M to 0.005 μ M in assays evaluating LPS- and FMLP-induced inhibition of tumor necrosis factor α (TNF- α) and leukotriene B4 (LTB4) in human whole blood. In a test measuring IgE-mediated allergic pulmonary inflammation induced by inhalation of antigen by sensitized guinea pigs, administration of I resulted in a significant reduction in the eosinophilia and the accumulation of other inflammatory leukocytes and effected less inflammatory lung damage. One hundred fifty-five invention compds. also inhibited the hydrolysis of cAMP to AMP by human recombinant phosphodiesterase IVa with IC50 values ranging from 160 nM to 0.086 nM. Thus, I and their pharmaceutical compns. are useful for the treatment or prevention of a variety of allergic, inflammatory, CNS, and other conditions (no data).

- IT 638220-30-1P, trans-2-[3'-(6-[1-(Methylsulfonyl)-1-methylethyl]quinolin-8-yl)biphenyl-4-yl]cyclopropanecarboxylic acid
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (PDE4 inhibitor; preparation of 8-arylquinoline PDE4 inhibitors for treatment of a variety of allergic, inflammatory, CNS, and other conditions)
 RN 638220-30-1 CA
 CN Cyclopropanecarboxylic acid, 2-[3'-(6-[1-methyl-1-(methylsulfonyl)ethyl]-8-quinolinyl)][1,1'-biphenyl]-4-yl]-, (1R,2R)-rel- (CA INDEX NAME)

Relative stereochemistry.



REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 4 OF 4 CA COPYRIGHT 2008 ACS on STN
 ACCESSION NUMBER: 138:73184 CA
 TITLE: Preparation of substituted 8-arylquinoline
 phosphodiesterase-4 (PDE4) inhibitors
 INVENTOR(S): Dube, Daniel; Girard, Yves; MacDonald, Dwight;
 Mastracchio, Anthony; Gallant, Michel; Lacombe,
 Patrick; Deschenes, Denis
 PATENT ASSIGNEE(S): Merck Frosst Canada & Co., Can.
 SOURCE: PCT Int. Appl., 204 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------------------|----------|-----------------|------------|
| WO 2003002118 | A1 | 20030109 | WO 2002-CA953 | 20020626 |
| W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW | | | | |
| RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG | | | | |
| CA 2450686 | A1 | 20030109 | CA 2002-2450686 | 20020626 |
| AU 2002344885 | A1 | 20030303 | AU 2002-344885 | 20020626 |
| AU 2002344885 | B2 | 20060629 | | |
| EP 1404330 | A1 | 20040407 | EP 2002-742600 | 20020626 |
| EP 1404330 | B1 | 20050601 | | |
| R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR | | | | |
| JP 2005501822 | T | 20050120 | JP 2003-508357 | 20020626 |
| AT 296630 | T | 20050615 | AT 2002-742600 | 20020626 |
| ES 2242036 | T3 | 20051101 | ES 2002-742600 | 20020626 |
| US 20040162314 | A1 | 20040819 | US 2003-478791 | 20031125 |
| US 6919353 | B2 | 20050719 | | |
| PRIORITY APPLN. INFO.: | | | US 2001-301220P | P 20010627 |
| | | | US 2001-303472P | P 20010706 |
| | | | WO 2002-CA953 | W 20020626 |
| OTHER SOURCE(S): | MARPAT 138:73184 | | | |
| GI | | | | |

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB 8-Arylquinolines (shown as I; variables defined below; e.g. both enantiomers of 4-hydroxy-1-[3-[6-(1-methanesulfonyl-1-methylethyl)quinolin-8-yl]phenyl]-2-(4-methanesulfonylphenyl)-4-methylpentan-3-one) wherein the aryl group at the 8-position contains a meta two atom bridge to an optionally substituted Ph or pyridyl group, are PDE4 inhibitors useful to treat asthma, chronic bronchitis, chronic obstructive pulmonary disease, arthritis, respiratory distress syndrome, allergic rhinitis, neurogenic

inflammation, pain, rheumatoid arthritis, and other diseases. R1-R7 and Ar are as in claim 1. For I: Ar is Ph, pyridinone, pyridyl, or pyridyl N-oxide, optionally substituted with 1-5 independent -Cl-6-alkyl, -OH, -CN, halogen, -CF₃, -(C0-6-alkyl)-SOn-(Cl-6-alkyl), -(C0-6-alkyl)-SOn-NH-(Cl-6-alkyl) or 5-membered heteroaryl ring containing 1-4 heteroatoms = O, S or N, wherein the 5-membered-ring is optionally substituted. R1 is H, halogen; or a -Cl-6-alkyl, -cycloC3-6alkyl, -Cl-6-alkenyl, -C0-4alkyl-C(O)-C0-4alkyl, -Cl-6-alkoxy, aryl, heteroaryl, -CN, -heterocycloC3-6-alkyl, -amino, -Cl-6-alkylamino, -(Cl-6-alkyl)(Cl-6-alkyl)amino, -Cl-6-alkyl(oxy)Cl-6-alkyl, -C(O)NH(aryl), -C(O)NH(heteroaryl), -SOnNH(aryl), -SOnNH(heteroaryl), -SOnNH(Cl-6-alkyl), -C(O)N(C0-6alkyl)(C0-6-alkyl), -NH-SOn-(Cl-6-alkyl), -carbamoyl, -(Cl-6-alkyl)-O-C(CN)dialkylamino, or -(C0-6-alkyl)-SOn-(Cl-6-alkyl) group, wherein any of the groups is optionally substituted with = 1-5 substituents. R2, R3, R6, and R7 = H, halogen, hydroxy, -Cl-6-alkyl, or -Cl-6-alkoxy, wherein the alkyl and alkoxy are optionally substituted independently with 1-3 halogen or OH; R4 is H, halogen, -CN, Ph, oxadiazolyl, or -C(O)-O-C0-6alkyl, wherein the alkyl and latter three possibilities are optionally substituted. R5 is H, hydroxy, -CN; or a -Cl-6-alkyl, -C(O)Cl-6alkyl, -C(O)aryl, -C(O)pyridyl, -C(O)-O-C0-6-alkyl, -C(O)-C3-7-cycloalkyl, -Cl-6-alkyl-C3-7cycloalkyl, -Cl-6-alkyl(C3-7-cycloalkyl)2, -Cl-6-alkylaryl, -C(O)-N(C0-6alkyl)2, -SOnaryl, -SOn-Cl-6-alkyl, -SOn-C3-7-cycloalkyl, -SOn-N(C0-6-alkyl)2, -P(O)(Cl-6-alkyl)2, -P(O)(Cl-6-alkoxy)2, Ph, pyridyl, -SOnimidazolyl, -SOnthiazolyl, 5-membered heteroaryl ring containing 1-4 heteroatoms = O, S or N or oxoisoxaphosphinanyl group, any of which group optionally substituted; or R5 and R6 form :O; or R6 and R3 form -CH2- or -O-; and n is 0-2. Although the methods of preparation are not claimed, >100 example preps. are included. The IC50 values for PDE4 inhibition of Examples 1-113 generally are 0.02-26 μ M as measured using LPS and FMLP-induced TNF- α and LTB4 assays in human whole blood. I were tested for effects on an IgE-mediated allergic pulmonary inflammation induced by inhalation of antigen by sensitized guinea pigs; Administration of I (0.001-10 mg/kg i.p. or p.o.), up to three times during the 48 h following antigen challenge, lead to a significant reduction in the eosinophilia and the accumulation of other inflammatory leukocytes. There was also less inflammatory damage in the lungs of animals treated with I. Comps. which inhibit the hydrolysis of cAMP to AMP by the type-IV cAMP-specific phosphodiesterases were screened in a 96-well plate format; IC50 values of I generally ranged 0.1-25 nM.

- IT 481680-95-9P, 2-[3-[6-(1-Methanesulfonyl-1-methylethyl)quinolin-8-yl]phenyl]-1-(4-methanesulfonylphenyl)cyclopropanecarboxylic acid
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (drug candidate; preparation of substituted 8-arylquinoline phosphodiesterase-4 (PDE4) inhibitors)
 RN 481680-95-9 CA
 CN Cyclopropanecarboxylic acid, 2-[3-[6-[1-methyl-1-(methylsulfonyl)ethyl]-8-quinolinyl]phenyl]-1-[4-(methylsulfonyl)phenyl]- (CA INDEX NAME)



REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> file marpat

=> s l1 full

L5 8 SEA SSS FUL L1

=> d ibib abs fqhit 1-8

L5 ANSWER 1 OF 8 MARPAT COPYRIGHT 2008 ACS on SIN


ACCESSION NUMBER: 148:472072 MARPAT
 TITLE: Substituted pyrazinone derivatives as selective 2C-adrenoceptor antagonists for use as a medicine and their preparation
 INVENTOR(S): Andres-Gil, Jose Ignacio; Alcazar-Vaca, Manuel Jesus; Linares De La Morena, Maria Lourdes; Martinez Gonzalez, Sonia; Oyarzabal Santamarina, Julien; Pastor-Fernandez, Joaquin; Vega-Ramiro, Juan Antonio; Delgado-Jimenez, Francisca; Drinkenburg, Wilhelmus Helena Ignatius Maria
 PATENT ASSIGNEE(S): Janssen Pharmaceutica NV, Belg.
 SOURCE: PCI Int. Appl., 74pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|--|------|----------|-----------------|----------|
| WO 2008043775 | A1 | 20080417 | WO 2007-EP60748 | 20071010 |
| W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, | | | | |

PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN,
 TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW
 RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,
 IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR, BF,
 BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW,
 GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ,
 BY, KG, KZ, MD, RU, TJ, TM


PRIORITY APPLN. INFO.:
 GI

EP 2006-122173 20061012




AB The invention concerns substituted pyrazinone derivs. according to the general formula I a pharmaceutically acceptable acid or base addition salt thereof, a stereochem. isomeric form thereof, an N-oxide form thereof or a quaternary ammonium salt thereof, having selective 2C-adrenoceptor antagonist activity. It further relates to their preparation, compns. comprising them and their use as a medicine. The compds. according to the invention are useful for the prevention and/or treatment of central nervous system disorders, mood disorders, anxiety disorders, stress-related disorders associated with depression and/or anxiety, cognitive disorders, personality disorders, schizoaffective disorders, Parkinson's disease, dementia of the Alzheimer's type, chronic pain conditions, neurodegenerative diseases, addiction disorders, mood disorders and sexual dysfunction. Compds. of formula I wherein A1 and A2 are independently N and C, with the proviso that not both A1 and A2 are C simultaneously; Z1, Z2 are independently a bond and NH and derivs.; n is 0, 1, 2, and 3; R5 is H and halo; P is Ph, biphenyl, 1,1-diphenylmethyl and benzyloxyphenyl; X2 is a bond, (un)saturated (un)substituted C1-8 hydrocarbon wherein one or more of the bivalent CH2 units may be replaced with bivalent Ph unit; Q2 is NH and derivs., nitrogen-heterocycle, OH and derivs., SH and derivs., SO2H and derivs., aryl, etc.; and their pharmaceutically acceptable acid and base addition salts, N-oxides and quaternary ammonium salts thereof, are claimed. Example compound II was prepared by N-alkylation of 3-(1-(biphenyl-4-ylmethyl)piperidin-4-yl)pyrazin-2-one with N-(2-chloroethyl)piperidine. All the invention compds. were evaluated for their 2C-adrenoceptor antagonistic activity (data given).


MSTR 1



G1 = 19-9 20-195



G2 = N
 G3 = bond
 G13 = 1-8 11-12



G14 = phenylene
 G18 = quinolinyl (opt. substd. by G42)
 G42 = Ph

Patent location:

claim 1

Note:

or pharmaceutically acceptable acid or base
 addition salts, N-oxides or quaternary ammonium
 salts

Note:

additional derivatization also claimed

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 2 OF 8 MARPAT COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 146:100694 MARPAT

TITLE: Preparation of piperidine derivatives as NMDA receptor antagonists

INVENTOR(S): Masui, Moriyasu; Matsumura, Akira

PATENT ASSIGNEE(S): Shionogi & Co., Ltd., Japan

SOURCE: PCT Int. Appl., 111pp.

CODEN: PIXXD2

DOCUMENT TYPE:

Patent

LANGUAGE:

Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|--|------|----------|------------------|----------|
| WO 2006137465 | A1 | 20061228 | WO 2006-JP312466 | 20060622 |
| W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, | | | | |

KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN,
 MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU,
 SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG,
 US, UZ, VC, VN, ZA, ZM, ZW
 RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,
 IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ,
 CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH,
 GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,
 KG, KZ, MD, RU, TJ, TM

PRIORITY APPLN. INFO.:

JP 2005-185100 20050624


JP 2005-309760 20051025

GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *


AB Title compds. I [A1 = nitrogenated aromatic monocyclic group or nitrogenated aromatic fused-ring group which has at least one (un)protected hydroxy and/or amino and which may be substituted by other group, nitrogenated aromatic monocyclic group or nitrogenated aromatic fused-ring group which has -NH- in the ring and in which other ring-constituting atom may have substituent(s) (except (un)protected hydroxy and amino); A2 = (un)substituted aromatic cyclic hydrocarbon, (un)substituted aromatic heterocycle; R1 = H, hydroxy, acyloxy, etc.; R2 = H, hydroxy, alkyl; R1 and R2 may combine to form single bond; m = 0, 1; X = (un)substituted alkenylene, (un)substituted alkenylene, -CO(CR3R4)n-, etc.; R3, R4 = H, (un)substituted alkyl; n = 0-4; when m is 0, Y represents single bond, -O-, -S-, etc.; when m is 1, Y represents single bond, alkylene, alkenylene, etc.], pharmaceutically acceptable salts or solvates thereof were prepared. For example, EDCI mediated amidation of 4-imidazolecarboxylic acid with compound II, e.g., prepared from 4-(4-chlorophenyl)-1,2,3,6-tetrahydropyridine hydrochloride in 2 steps, afforded compound III [R = imidazol-4-yl] in 35% yield. In NMDA receptor (NR1/NR2B receptor) binding assays, the IC50 value of compound III [R = 2,3-dihydro-2-oxo-1H-benzimidazol-5-yl] was 0.002 μ M. Compds. I are claimed useful as analgesics.

MSTR 1



G1 = quinolinyl (subst'd. by OH)

G6 = 349



G9 = phenylene
 G22 = bond
 G25 = 3-2 4-5 17-143




Patent location: claim 1
 Note: substitution is restricted
 Note: or pharmaceutically acceptable salts or solvates

REFERENCE COUNT: 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 3 OF 8 MARPAT COPYRIGHT 2008 ACS on SIN
 ACCESSION NUMBER: 144:610 MARPAT
 TITLE: Phenol analogs for inducing hematopoietic stem cells and megakaryocytes
 INVENTOR(S): Kashikura, Ikuo; Yoshizawa, Atsushi
 PATENT ASSIGNEE(S): Ohtsuka Pharmaceutical Co., Ltd., Japan
 SOURCE: Jpn. Kokai Tokkyo Koho, 16 pp.
 CODEN: JKXXAF
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------------------|------|----------|-----------------|----------|
| JP 2005330204 | A | 20051202 | JP 2004-148779 | 20040519 |
| PRIORITY APPLN. INFO.: | | | JP 2004-148779 | 20040519 |

GI



AB Phenol analogs (I; R1 = OH, Me, alkyl, alkoxy; R2 = (substituted) aryl or heteroaryl; R3 = H, halogen, alkyl, etc.; n = 2-4; A1, A2 = N, CH, C(R3)), including 4-(5-octylpyrimidin-2-yl)phenol, 4-[2-(4-butoxyphenyl)pyrimidin-5-yl]phenol, 2-(octylphenyl)pyrimidin-5-ol, 4-(5-octylpyrimidin-2-yl)phenol, 4-(6-pentylquinolin-2-yl)phenol or 4-[2-[4-(disiloxy)-2,3-difluorophenyl]pyrimidin-5-yl]phenol., are claimed for inducing hematopoietic stem cells and megakaryocytes for treatment of platelet diseases, anemia, leukemia, SLE, DIC, tumor, etc.


MSTR 1



G2 = (1) G5
G3 = 11



G5 = quinolinyl (opt. substd. by alkyl <containing 1-20 C>) / 27



Patent location: claim 1
Note: or salts

L5 ANSWER 4 OF 8 MARPAT COPYRIGHT 2008 ACS on STN
ACCESSION NUMBER: 141:410960 MARPAT
TITLE: Preparation of 8-(3-biaryl)phenylquinoline
phosphodiesterase-4 inhibitors
INVENTOR(S): Dube, Daniel; Dube, Laurence; Gallant, Michel;
Lacombe, Patrick; Deschenes, Denis; MacDonald, Dwight
PATENT ASSIGNEE(S): Merck Frosst Canada & Co., Can.
SOURCE: PCT Int. Appl., 129 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------|----------|-----------------|----------|
| WO 2004096220 | A1 | 20041111 | WO 2004-CA622 | 20040427 |
| W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, | | | | |

SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE,
 SN, TD, TG


| | | | | |
|--|----|----------|------------------|----------|
| AU 2004234190 | A1 | 20041111 | AU 2004-234190 | 20040427 |
| CA 2523336 | A1 | 20041111 | CA 2004-2523336 | 20040427 |
| EP 1635829 | A1 | 20060322 | EP 2004-729586 | 20040427 |
| R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK | | | | |
| CN 1812787 | A | 20060802 | CN 2004-80018346 | 20040427 |
| JP 2006524638 | T | 20061102 | JP 2006-504121 | 20040427 |
| US 20060223850 | A1 | 20061005 | US 2005-554176 | 20051021 |
| IN 2005DN04934 | A | 20070928 | IN 2005-DN4934 | 20051027 |
| PRIORITY APPLN. INFO.: | | | US 2003-466542P | 20030430 |
| | | | WO 2004-CA622 | 20040427 |

GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The title 8-phenylquinolines I [S1-S3 = H, OH, halo, alkyl, etc.; R1 = CO2aryl, CONHaryl, CONHheteroaryl, etc.; Ar1, Ar2 = (hetero)aryl or an N-oxide thereof; R2 = H, aryl, haloaryl, heterocyclyl, etc.; R3 = H, alkyl, hydroxyalkyl, etc.; R4 = H, halo, CN, alkyl, etc.] which are PDE4 inhibitors, were prepared E.g., a multi-step synthesis of II (no characterization data given for intermediates), which showed IC50 of 0.155 µM in LPS and FMLP-induced TNF-α and LTB4 assays in human whole blood, was given. The pharmaceutical compns. comprising the compound I are claimed.

MSTR 1



G4 = carbocycle <containing 3-6 C, 1-3 rings>
 G16 = p-C6H4
 G33 = 92-18 91-53 94-52



Patent location: claim 1
 Note: or pharmaceutically acceptable salts

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 5 OF 8 MARPAT COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 140:59526 MARPAT
 TITLE: Preparation of 8-(biaryl)quinolines as PDE4 inhibitors
 INVENTOR(S): Deschenes, Denis; Dube, Daniel; Dube, Laurence;
 Gallant, Michel; Girard, Yves; Lacombe, Patrick;
 MacDonald, Dwight

PATENT ASSIGNEE(S): Merck Frosst Canada & Co., Can.
 SOURCE: PCT Int. Appl., 122 pp.
 CODEN: PIXXD2

DOCUMENT TYPE: Patent
 LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------------------|--|----------|-----------------|----------|
| WO 2004000814 | A1 | 20031231 | WO 2003-CA957 | 20030623 |
| W: | AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW | | | |
| RW: | GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG | | | |
| CA 2490043 | A1 | 20031231 | CA 2003-2490043 | 20030623 |
| AU 2003243870 | A1 | 20040106 | AU 2003-243870 | 20030623 |
| EP 1517895 | A1 | 20050330 | EP 2003-760540 | 20030623 |
| EP 1517895 | B1 | 20070314 | | |
| R: | AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK | | | |
| JP 2006502104 | T | 20060119 | JP 2004-514482 | 20030623 |
| AT 356808 | T | 20070415 | AT 2003-760540 | 20030623 |
| ES 2282667 | T3 | 20071016 | ES 2003-760540 | 20030623 |
| US 20050234238 | A1 | 20051020 | US 2004-517416 | 20041208 |
| US 7153968 | B2 | 20061226 | | |
| PRIORITY APPLN. INFO.: | | | US 2002-391364P | 20020625 |
| | | | US 2002-428313P | 20021122 |
| | | | WO 2003-CA957 | 20030623 |

GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Title compds. I [wherein A = C or N; X = Ph, pyridyl, pyrazinyl, thiaphenyl, quinolinyl, benzofuranyl, oxadiazolyl, diazolyipyridinyl, imidazolypyridinyl, oxadiazolylphenyl, benzodioxolyl; R1 = H, halo, or (un)substituted alkanoyl, cyclo/alkyl, alkenyl; R2, R3 = independently H, halo, OH, CN, NO2, or dialkenyl/dicycloalkyl/alkyl, alkenyl, wide variety of C-containing and heteroat. groups and/or functional groups optionally linked by C1-4alkyl; R2 optionally forms a double bond with an adjoining bond; R4 = H, halo; any ring nitrogen optionally forms N-oxide and N-chloride; and pharmaceutically acceptable salts thereof] were prepared as phosphodiesterase IV (PDE4) inhibitors. For example, II was prepared by Suzuki cross-coupling of quinoline III with 2-bromo-3-chlorothiophene. One hundred fifty-five invention compds. suppressed PDE4 with IC50 values ranging from 36 μ M to 0.005 μ M in assays evaluating LPS- and FMLP-induced inhibition of tumor necrosis factor α (TNF- α) and leukotriene B4 (LTB4) in human whole blood. In a test measuring IgE-mediated allergic pulmonary inflammation induced by inhalation of antigen by sensitized guinea pigs, administration of I resulted in a significant reduction in the eosinophilia and the accumulation of other inflammatory leukocytes and effected less inflammatory lung damage. One hundred fifty-five invention compds. also inhibited the hydrolysis of cAMP to AMP by human recombinant phosphodiesterase IVa with IC50 values ranging from 160 nM to 0.086 nM. Thus, I and their pharmaceutical compns. are useful for the treatment or prevention of a variety of allergic, inflammatory, CNS, and other conditions (no data).

MSTR 1



G1 = 19

C-G2
19

G3 = 86

P-C₈H₄G13

G5 = cycloalkyl <containing 3-6 C>

(opt. substd. by (1-6) G6)
G13 = 129



Patent location: claim 1
Note: or pharmaceutically acceptable salts, N-oxides or N-chlorides

REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 6 OF 8 MARPAT COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 138:73184 MARPAT

TITLE: Preparation of substituted 8-arylquinoline phosphodiesterase-4 (PDE4) inhibitors

INVENTOR(S): Dube, Daniel; Girard, Yves; MacDonald, Dwight; Mastracchio, Anthony; Gallant, Michel; Lacombe, Patrick; Deschenes, Denis

PATENT ASSIGNEE(S): Merck Frosst Canada & Co., Can.

SOURCE: PCT Int. Appl., 204 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:


| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|------------------------|--|----------|-----------------|----------|
| WO 2003002118 | A1 | 20030109 | WO 2002-CA953 | 20020626 |
| W: | AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW | | | |
| RW: | GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG | | | |
| CA 2450686 | A1 | 20030109 | CA 2002-2450686 | 20020626 |
| AU 2002344885 | A1 | 20030303 | AU 2002-344885 | 20020626 |
| AU 2002344885 | B2 | 20060629 | | |
| EP 1404330 | A1 | 20040407 | EP 2002-742600 | 20020626 |
| EP 1404330 | B1 | 20050601 | | |
| R: | AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR | | | |
| JP 2005501822 | T | 20050120 | JP 2003-508357 | 20020626 |
| AT 296630 | T | 20050615 | AT 2002-742600 | 20020626 |
| ES 2242036 | T3 | 20051101 | ES 2002-742600 | 20020626 |
| US 20040162314 | A1 | 20040819 | US 2003-478791 | 20031125 |
| US 6919353 | B2 | 20050719 | | |
| PRIORITY APPLN. INFO.: | | | US 2001-301220P | 20010627 |
| | | | US 2001-303472P | 20010706 |
| | | | WO 2002-CA953 | 20020626 |

GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB 8-Arylquinolines (shown as I; variables defined below; e.g. both enantiomers of 4-hydroxy-1-[3-(6-(1-methanesulfonyl-1-methylethyl)quinolin-8-yl)phenyl]-2-(4-methanesulfonylphenyl)-4-methylpentan-3-one) wherein the aryl group at the 8-position contains a meta two atom bridge to an optionally substituted Ph or pyridyl group, are PDE4 inhibitors useful to treat asthma, chronic bronchitis, chronic obstructive pulmonary disease, arthritis, respiratory distress syndrome, allergic rhinitis, neurogenic inflammation, pain, rheumatoid arthritis, and other diseases. R1-R7 and Ar are as in claim 1. For I: Ar is Ph, pyridinone, pyridyl, or pyridyl N-oxide, optionally substituted with 1-5 independent -C1-6-alkyl, -OH, -CN, halogen, -CF₃, -(C0-6-alkyl)-SOn-(C1-6-alkyl), -(C0-6-alkyl)-SOn-NH-(C1-6-alkyl) or 5-membered heteroaryl ring containing 1-4 heteroatoms = O, S or N, wherein the 5-membered-ring is optionally substituted. R1 is H, halogen; or a -C1-6-alkyl, -cycloC3-6alkyl, -C1-6-alkenyl, -C0-4alkyl-C(O)-C0-4alkyl, -C1-6-alkoxy, aryl, heteroaryl, -CN, -heterocycloC3-6-alkyl, -amino, -C1-6-alkylamino, -(C1-6-alkyl)(C1-6-alkyl)amino, -C1-6-alkyl(oxy)C1-6-alkyl, -C(O)NH(aryl), -C(O)NH(heteroaryl), -SOnNH(aryl), -SOnNH(heteroaryl), -SOnNH(C1-6-alkyl), -C(O)N(C0-6alkyl)(C0-6-alkyl), -NH-SOn-(C1-6-alkyl), -carbamoxy, -(C1-6-alkyl)-O-C(N)alkylamino, or -(C0-6-alkyl)-SOn-(C1-6-alkyl) group, wherein any of the groups is optionally substituted with = 1-5 substituents. R2, R3, R6, and R7 = H, halogen, hydroxy, -C1-6-alkyl, or -C1-6-alkoxy, wherein the alkyl and alkoxy are optionally substituted independently with 1-3 halogen or OH; R4 is H, halogen, -CN, Ph, oxadiazolyl, or -C(O)-O-C0-6alkyl, wherein the alkyl and latter three possibilities are optionally substituted. R5 is H, hydroxy, -CN; or a -C1-6-alkyl, -C(O)C1-6alkyl, -C(O)aryl, -C(O)pyridyl, -C(O)-O-C0-6-alkyl, -C(O)-C3-7-cycloalkyl, -C1-6-alkyl-C3-7cycloalkyl, -C1-6-alkyl(C3-7-cycloalkyl)2, -C1-6-alkylaryl, -C(O)-N(C0-6alkyl)2, -SOnaryl, -SOn-C1-6-alkyl, -SOn-C3-7-cycloalkyl, -SOn-N(C0-6-alkyl)2, -P(O)(C1-6-alkyl)2, -P(O)(C1-6-alkoxy)2, Ph, pyridyl, -SOnimidazolyl, -SOnthiazolyl, 5-membered heteroaryl ring containing 1-4 heteroatoms = O, S or N or oxoloxaphosphinanyl group, any of which group optionally substituted; or R5 and R6 form -O-; or R6 and R3 form -CH2- or -O-, and n is 0-2. Although the methods of preparation are not claimed, >100 example preps. are included. The IC50 values for PDE4 inhibition of Examples 1-113 generally are 0.02-26 μM as measured using LPS and FMLP-induced TNF-α and LTB4 assays in human whole blood. I were tested for effects on an IgE-mediated allergic pulmonary inflammation induced by inhalation of antigen by sensitized guinea pigs;. Administration of I (0.001-10 mg/kg i.p. or p.o.), up to three times during the 48 h following antigen challenge, lead to a significant reduction in the eosinophilia and the accumulation of other inflammatory leukocytes. There was also less inflammatory damage in the lungs of animals treated with I. Compds. which inhibit the hydrolysis of cAMP to AMP by the type-IV cAMP-specific phosphodiesterases were screened in a 96-well plate format; IC50 values of I generally ranged 0.1-25 nM.

MSTR 1



G1 = carbocycle <containing 3-6 C, mono- or polycyclic>
(opt. substd.)
G4 = Ph (opt. substd. by (1-3) G15)
G30 = 196-13 197-22



G31 = O
Patent location: claim 1
Note: or pharmaceutically acceptable salts

REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 7 OF 8 MARPAT COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 135:288703 MARPAT
TITLE: 3-Cyanoquinolines, 3-cyano-1,6-naphthyridines, and 3-cyano-1,7-naphthyridines as protein kinase inhibitors
INVENTOR(S): Boschelli, Diane Harris; Wang, Yanong; Boschelli, Frank Charles; Berger, Dan Maarten; Zhang, Nan; Powell, Dennis William; Ye, Fei; Yamashita, Ayako; Demorin, Frenel Fils; Wu, Biqi; Tsou, Hwei-ru; Overbeek-klumpers, Elsebe Geraldine; Wissner, Allan
PATENT ASSIGNEE(S): American Home Products Corporation, USA
SOURCE: PCT Int. Appl., 448 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|--|------|----------|-----------------|----------|
| WO 2001072711 | A1 | 20011004 | WO 2001-US9966 | 20010328 |
| W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, | | | | |

RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN,
 YU, ZA, ZW
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY,
 DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF,
 BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
 CA 2404662 A1 20011004 CA 2001-2404662 20010328
 EP 1268431 A1 20030102 EP 2001-924407 20010328
 EP 1268431 B1 20070829
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
 IE, SI, LT, LV, FI, RO, MK, CY, AL, TR
 BR 2001009650 A 20030422 BR 2001-9650 20010328
 JP 2003528857 T 20030930 JP 2001-570624 20010328
 AT 371647 T 20070915 AT 2001-924407 20010328
 ES 2291311 T3 20080301 ES 2001-924407 20010328
 MX 2002PA09439 A 20030224 MX 2002-PA9439 20020926
 PRIORITY APPLN. INFO.: US 2000-535843 20000328
 WO 2001-US9966 20010328

GI


* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Title compds. I [X = N(H) or substituted derivs., O, SOO-2; n = 0-1; A =
 divalent (un)substituted alkyl, C(O), C(O)-alkyl, alkyl-C(O), cycloalkyl,
 or absent; T, Z = C, N provided that both T and Z are not N; R1 =
 cycloalkyl, 5-6 atom (hetero)aryl ring containing 0-4 heteroatoms, 8-20 atom
 bicyclic heteroaryl ring containing 1-4 heteroatoms, etc.; R2a-c = H, aryl,
 CH2-aryl, O-aryl, SOO-2-aryl, NO2, SH, etc.; R3 = alkenyl, alkynyl,
 (hetero)aryl; R4 = (un)substituted alkyl, alkenyl, alkynyl, (hetero)aryl]
 were prepared. Over 500 synthetic examples were disclosed, including some
 combinatorial preps., and addnl. reference examples. E.g.,
 4-[(4-bromo-2-thienyl)methyl]morpholine reacted with
 bis(pinacolato)diboron [DMSO, PdCl2(dppf), KOAc] to give dioxaborolane II.
 II was coupled to 7-bromo-4-[3-chloro-4-[(1-methyl-1H-imidazol-2-
 yl)sulfanyl]anilino]-3-quinolinecarbonitrile [preparation given; diglyme,
 Pd(PPh3)4, NaHCO3] to yield invention compound III as a yellow solid after
 purification. III had IC50 = 6.0 nM for Raf1 kinase and inhibited the human
 adenocarcinoma CaCo-2 cell line with IC50 = 1.9, 0.78 (2 trials). I are
 useful as antineoplastic agents, and in the treatment of osteoporosis and
 polycystic kidney disease.

MSTR 1

G26-G1-G6-G9

G1 = 36-1 40-3



G3 = 53-38 54-36




G4 = 261



G25 = OMe
G26 = 274



G34 = 282



G35 = (0-5) CH2
G36 = 285



G38 = pyrrolidino
G50 = 422-2 424-275



G51 = CH
Patent location:
Note:

claim 1
substitution is restricted

Note: or pharmaceutically acceptable salts

REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 8 OF 8 MARPAT COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 125:142750 MARPAT
 TITLE: Polyarylcabamoylaza- and -carbamoylalkanedioic acids as squalene synthase inhibitors

INVENTOR(S): Pauls, Henry W.; Choi, Yong-Mi; Studt, Robert W.; Maguire, Martin P.; Spada, Alfred P.; Cha, Don D.

PATENT ASSIGNEE(S): Rhone-Poulenc Rorer Pharmaceuticals Inc., USA
 SOURCE: PCT Int. Appl., 55 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent



LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

| PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|---|------|----------|-----------------|----------|
| WO 9618615 | A1 | 19960620 | WO 1995-US15364 | 19951129 |
| W: AL, AM, AT, AU, BB, BG, BR, BY, CA, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ | | | | |
| RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG | | | | |
| US 5556990 | A | 19960917 | US 1994-357481 | 19941216 |
| CA 2207429 | A1 | 19960620 | CA 1995-2207429 | 19951129 |
| AU 9643698 | A | 19960703 | AU 1996-43698 | 19951129 |
| AU 695852 | B2 | 19980827 | | |
| EP 801644 | A1 | 19971022 | EP 1995-942489 | 19951129 |
| R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE | | | | |
| JP 10511084 | T | 19981027 | JP 1995-518973 | 19951129 |
| PRIORITY APPLN. INFO.: | | | US 1994-357481 | 19941216 |
| | | | WO 1995-US15364 | 19951129 |

GI




II

AB This invention relates to a class of novel dicarboxy amide derivs. of lipophilic amines I wherein: A is O, S, NR, SO, SO₂, or a bond; B is (CRR)₁₋₂, O, S, NR, SO, SO₂, RC:CR, C.tplbond.C, CO, or a bond; Y is, e.g., RNZ(CRR)dCRR, N-Z-piperidyl, where Z is COWCR7[(CR3R4)fCO2R][(CR5R6)gCO2R]; W is a bond, (CRR)h, or NR; R = H, alkyl; R₁, R₂ are independently H, alkyl, alkoxy, OH, halo, haloalkyl, Ph; R₃-R₆ are independently H, alkyl; R₇ is H, NRR, or OH and when W is (CRR)h then R₇ is OH; one of R₃-R₇ is OH; Ar₁ and Ar₂ are independently a mono- or diaryl or heteroaryl; p and q are independently 0-3; p + q is 0-4; d is 0-3; p + q + d is 1-3; f is 0-2; g is 0-2; h is 1-2; m and n are independently 0-2; which exhibit squalene synthase inhibition properties. Comps. of this invention reduce levels of serum cholesterol in the body without significantly reducing mevalonic metabolite synthesis. This invention relates also to pharmacol. compns. and method of treatment for lowering serum cholesterol levels using the compds. of this invention. Thus, e.g., coupling of prepared intermediates 3-hydroxy-3-(4-naphth-2-ylphenyl)piperidine with 3-hydroxy-3,4-bis(ethoxycarbonyl)butanoic acid afforded the diester intermediate which was hydrolyzed to the diaryl carbamoyl alkanedioic acid II which exhibited inhibition of squalene synthase with IC₅₀ = 27 nM.

MSTR 1B

G1—G16—G18

G1 = 82



G12 = Ph
G16 = phenylene (opt. substd. by (1-2) G12)
G18 = quinolinyl (opt. substd. by (1-2) G12)
G20 = Ph
Derivative: or pharmaceutically acceptable salts
Patent location: claim 1
Note: substitution is restricted
Stereochemistry: stereoisomers, enantiomers, diastereoisomers, and
racemic mixtures

=> d his

(FILE 'HOME' ENTERED AT 15:36:41 ON 31 JUL 2008)

FILE 'REGISTRY' ENTERED AT 15:37:58 ON 31 JUL 2008

L1 STRUCTURE UPLOADED

L2 13 S L1 SAM

L3 201 S L1 FULL

FILE 'CA' ENTERED AT 15:38:33 ON 31 JUL 2008

L4 4 S L3

FILE 'MARPAT' ENTERED AT 15:38:45 ON 31 JUL 2008

L5 8 S L1 FULL

=>

---Logging off of STN---

=>

Executing the logoff script...

=> LOG Y

STN INTERNATIONAL LOGOFF AT 15:40:13 ON 31 JUL 2008